

Dear New Patient:

To help treat your condition, your doctor has recommended a type of physical therapy called <u>Pelvic Floor</u> <u>Physical Therapy</u>. If you are unfamiliar with this type of therapy, you are not alone. However, research points to its high success rate, all without drugs or surgery. In fact, over 90% of our patients report an 80-100% improvement in their symptoms by the end of therapy, and only 1-2% report no change at all. So, the odds are very good that we will be able to help you.

Many bowel, bladder, and pain conditions in the pelvis are due to poorly functioning muscles. Your "pelvic floor" is a muscle group at the bottom of your pelvis that has two main functions: 1) postural support to the pelvic organs (bladder, bowel, uterus, and prostate) and 2) voluntary control over bowel and bladder function.

Pelvic Floor Muscle Dysfunction Categories

- 1. Weak and sagging
 - Loss of control over your bladder or bowels (incontinence)
 - Pelvic organ prolapse (fallen bladder, bowels, and/or uterus)
- 2. Tight and spasming
 - Trouble with elimination (urinary retention, constipation)
 - Pelvic pain (with sitting, during intercourse)

Pelvic Floor Rehab is <u>not</u> the same as Kegel exercises. Like traditional physical therapy, we will use techniques that will strengthen weak muscles and stretch and massage tight muscles. Our team has expert training in how to modify traditional therapy techniques in order to treat the pelvic floor area in ways that will make you feel comfortable.

Please take the time to fill out the enclosed questionnaire. It will help us better plan your treatment. There may be questions that at first, you wouldn't think would apply to you, but since pelvic floor problems can affect bowel, bladder and sexual function, it is not uncommon to have symptoms in more than one category.

Your evaluation and first treatment with your therapist will take about an hour, will be one-on one, and in a private, quiet area. Please dress in comfortable, loose-fitting clothing. You are welcome to bring a friend or family member with you. Most patients will require 12-16 treatment sessions that will last about 45-60 minutes each. Please call with any questions or concerns you might have. We look forward to meeting you.

Sincerely,

Your Pelvic Floor Rehab Team

Tracey Goldstein-Marquez, PT, DPT

Filamae Garnica-Tapiculin, PT

Nadine Gibson, PT

Lori Yoder, PTA

Ashley Clark, PTA



Male Pelvic Floor Questionnaire

HOW DID YOU HEAR ABOUT US? □ PHYSICIAN □ WEBSITE □ FACEBOOK □ SEMINAR □ NEWSPAPER AD □ FRIEND □ RETURNING PATIENT □ OTHER_____

Have you recently traveled out of this country? Yes □ No □ Have you had direct/indirect prolonged contact with someone with a confirmed case of coronavirus? Yes □ No □

Conditions related to the pelvic floor muscles can affect bladder, bowel, sexual, and physical function. Please answer the questions below so that we can better understand your problem. Please check all that apply.

History of Current Condition

Reason for today's visit

How long have you had this problem?

Since the problem began, has the problem become: D Worse D Better D Unchanged	
Is it related to an injury or accident? 🛛 No 🖵 Yes Explain how and when	
What are your goals for treatment?	

Previous <u>Treatments</u> for your condition:

Prostatectomy	Kegel exercises	Hemorrhoid repair	Massage therapy
□ Radiation	Biofeedback	Ostomy pouch	Chiropractor
Seed implantation	Pelvic floor rehab	Hemorrhoid cream	□ Acupuncture
Cyrotherapy	Diet/fluid changes	High fiber diet	Over the counter pain meds
Proton therapy	Bladder control meds	Probiotics	Prescription pain meds
Lupron/hormones	Collagen injections	Stool softeners	Bladder instillations
Greenlight ablation	□ Self-catheterization	Constipation meds	Urethral/bladder dilation
□ TURP	□ InterStim	Anti-diarrheal meds	Bladder instillations
□ BCG treatments	Artificial sphincter	Herbal supplements	Nerve injections/blocks
Condom catheter	Penile clamp	Enemas	Other:

Have you received therapy for the current or other problem in the past year? \Box Yes \Box No If yes, indicate type (physical therapy, speech therapy, etc.) and date_____

Previous <u>Tests</u> for your condition:

Urodynamics study	□ Video defacography	🗖 X-ray	Digital rectal exam
□ Bladder scan (PVR)	Colonoscopy	CT scan	Prostate biopsy
Cystoscopy	□ Anal manometry	□ MRI	Exploratory surgery

Medical Conditions and Health Status

Please rate your overall health:
Excellent
Good
Fair
Poor
With whom do you live?
Alone
Spouse/significant other
Other relative(s)
Roommate(s)
Where do you live?
Private home or apartment
Independent living
Assisted living
Other_____

	□ Enlarged prostate (BPH)	Prostate infections	Chronic prostatitis	Prostate pain					
	Erectile dysfunction	Erectile/ejaculation p	pain						
te	Prostate cancer: If so, when were you diagnosed? What was your Gleason score?								
state	Last PSA reading:								
Pro	Sexual Function: I'm sexually active								
1	☐ I'm not sexually active due to: □ My pelvic pain symptoms □ My other medical problems								
		For non-health relation	ated reasons D My part	tner's medical problem	IS				
	Urinary	Small bladder capacity	y 🛛 Bladder car	ncer	Bladder infections				
Bladder	incontinence	Large bladder capacity	y Interstitial	cystitis	(UTI's)				
ad	Overactive bladder	Kidney stones	□ Other		Painful bladder				
B	Urinary retention				syndrome				



Patient Name_____

Date

s	Fecal incontinenceConstipation		lcerative colitis pastic colon	HemorrhoidsLactose intolerance		Colon cancerAnal cancer		
Bowels	Diarrhea		verticulitis	Gluten intolerance		Colostomy/		
Bc	□ Irritable bowel		viverticulosis	• Other		ileostomy		
	syndrome							
	High blood pressureLow blood pressure		Asthma Cigarette smoker	DiabetesAcid reflux		Sciatica Stenosis		
	☐ High cholesterol		Former smoker	Depression		Arthritis		
	Heart disease		Migraine headaches	 Anxiety disorder 	-	Area		
Γ	Heart attack		Stroke	Glaucoma		Herniated disc		
Other Medical	Angina		TIA (mini strokes)	□ Cataracts		Level		
Ied	Congestive heart failur		Parkinson's Disease	Glasses		Degenerative disc		
r N	Cardiac arrhythmia		Multiple Sclerosis	Hearing loss		Area		
he	Decemaker/defibrillator		Seizures	Fibromyalgia		Bone fracture		
0	Vascular disease		Hypothyroidism	Osteoporosis		Area		
	Swollen legs/edema		Kidney disease	Osteopenia		Cancer		
	Emphysema		Kidney stones	Low back pain		Area		
	Bronchitis		Sleep apnea	Tailbone trauma		Other		
	Surgeries		Age or Year	Medications		For what condition?		
All	ergies							
נ 🗖	None 🛛 Latex sensitivit	y 🗆 S	Seasonal (pollen/hay fe	ver) 🗆 Bees 🗖 Other				
	Food		· ·	_ Ó				
Me	dications							
	view of Systems - Please ch			· · ·				
		Eye pai		Stomach pain		Tremors		
	Weight change	Chest p		Bloody stools		Speech changes		
	Fatigue/night sweats	D Poundi		□ Tarry stools		Seizures		
	Skin rash/itch	□ Shortne	ess of breath	Blood in urine		Loss of consciousness		
	Headaches	Calf pa	ain	Muscle pain	• •	Vertigo/spinning		
	Hearing loss	Leg sw	velling	Neck/back pain	٦U	Unsteadiness		
	Ringing in ears	Cough		Joint pain		Lightheaded		
	Ear pain/discharge	🛛 Coughi	ing up blood	□ Falls		Depression		
	Nose bleeds	D Phlegm	n production	□ Arm/leg weakness	ו 🗖	Nervous/anxious		
	Sore throat	Conges	stion	Lack of coordination		Suicidal thoughts		
	Blurred vision	U Wheez		Difficulty walking		Hallucinations		
	Double vision	Heartbu	urn	Difficulty standing up		Substance abuse		
	Light sensitive	Nausea	a/vomiting	Tingling/numbness		Memory loss		
	Is your doctor aware of these recent symptoms? Yes No							



Date_

Please answer the questions below. If your symptom severity fluctuates, characterize your symptoms at their worst.

Bladder Health Do you have a urologist? Yes No If yes, who?_____

- 1. During the <u>daytime</u>, how often do urinate? □ Every 30-60 minutes □ 1-2 hours □ 2-3 hours □ 3-4 hours □ More than 4 hours
- During the <u>nighttime</u> (after you've fallen asleep), how often do you get up to urinate? □ 0-1 times per night □ 2-3 times □ 3-4 times □ More than 4 times
- 3. How many 8 ounce servings (cups) do you drink of the following?

Liquid	Per Day	Per Week	On Occasion	Never	Liquid	Per Day	Per Week	On Occasion	Never
Coffee (Reg. Decaf)					Water				
Tea (Reg. Decaf)					Milk				
Soda (Reg. Decaf)					Juice				
Beer/Wine/Liquor					Other				

4. Do you every lose urine (even a few drops) with any of the situations below?

	Never	On Occasion	Sometimes	Usually	Always
Cough					
Sneeze					
Laugh					
Getting out of a chair					
Getting out of a car					
Getting out of bed at night					
Getting out of bed in the morning					
Picking objects off the floor					
Putting on shoes or clothes					
Lifting light items					
Lifting heavy items					
Cooking or doing dishes					
Doing housework/chores					
Doing yard work					
During sexual activity					
Walking to the toilet at home					
Shopping or running errands					
Standing or walking for a long time					
Walking to toilet in public					
Recreational activities					
Exercise activities					
Other					

5. Do you ever have strong or difficult-to-control urges to urinate with the situations listed below?

	Never	On occasion	Sometimes	Usually	Always
Around running water					
Feeling cold					
Feeling anxious or nervous					
Unlocking the door					
If I've waited too long					
Rushing off to the toilet					



- How long can you <u>usually delay an urge</u> to urinate? □ I rarely feel urges to void □ I go as soon as I feel an urge
 □ 1-2 minutes □ Several minutes □ 10-15 minutes □ 15 minutes or more
- 7. What <u>type of protective padding</u> do you use for bladder control? □ None needed □ Change underwear □ Folded tissue paper □ Liners □ Thin pads □ Thick pads □ Diapers □ Other_____
- 8. How often do you <u>change your bladder protection</u>? □ None □ Only when I leave the house □ Only at night
 □ Only during a cold □ Only during exercise □ 1-2 per day □ 3-4 per day □ 4+ per day
- 9. How<u>saturated</u> does your protection get? □ No leakage □ "Near misses" □ A few drops □ Damp □ Wet □ Soaked □ Overflows onto clothes
- 10. How often do you go to the bathroom <u>before you feel urges</u> to void, "just in case?" □ Never □ On occasion □ Sometimes □ Usually □ Always
- 11. How often do you <u>avoid drinking</u> fluid in order to help with bladder control? □ Never □ On occasion □ Sometimes □ Usually □ Always
- 12. Do you ever notice any of the following **<u>bladder symptoms</u>**? Please check how often.

	Never	On occasion	Sometimes	Usually	Always	
Weak stream						
Incomplete bladder emptying						
Trouble starting urine stream						
Strain to urinate						
Dribble after urinating						
Have to rock pelvis to empty bladder						
Have to push over the bladder to empty						
Splint or support bladder to urinate						
Pain as my bladder <i>fills</i>						
Location:	Bladder DU	rethra 🛛 Abdom	en	•	-	
Circle severity:	No pain 0 1 2	3 4 5 6 7 8	8 9 10 Worst pa	in		
Pain as my bladder empties						
Location:	Bladder Urethra Abdomen					
Circle severity:	No pain 0 1 2	3 4 5 6 7 8	8 9 10 Worst pa	uin		

Bowel Health Do you have a gastroenterologist? Yes No If yes, who?_

- 1. How often do you have a bowel movement? □ Less than 3 times a week □ Every 2-3 days □ Every 1-2 days □ Daily □ 2-3 times per day □ More than 3 times per day □ I won't go for several days, and then go multiple times in one day
- 2. What is the <u>consistency</u> of your bowel movements? □ Watery/formless □ Loose and thin □ Soft and formed □ Hard and rocky □ Small and pellet-like □ It varies_____
- 3. Do you ever <u>lose feces</u> with any of the situations below? Please check how often.

	Never	On Occasion	Sometimes	Usually	Always
On the way to the toilet					
If I exert myself					
When I pass gas					
I have fecal soiling without an urge to have a BM					

- 4. What <u>type of protective padding</u> do you use for bowel control? □ None needed □ Change underwear □ Folded tissue paper □ Liners □ Thin pads □ Thick pads □ Diapers □ Other_____
- 5. How often do you <u>change your bowel protection</u>? □ None □ Only when I leave the house □ Only when I have diarrhea/ loose stools □ 1-2 per day □ 3-4 per day □ 4+ per day



6. Do you every notice any of the following **<u>bowel symptoms</u>**? Please check how often.

	Never	On occasion	Sometimes	Usually	Always		
Excessive straining during a BM							
Support/splint rectum during BM							
Incomplete BM's							
Rush to the toilet with BM urge							
Trouble controlling gas in public							
Excessive wiping needed after BM							
Fecal soiling in underwear after BM							
Pain as my bowels <i>fill</i>							
Location:	Rectum Ab	domen					
Circle severity:	No pain 0 1 2	3 4 5 6 7 8	9 10 Worst pai	n			
Pain as my bowels <i>empty</i>							
Location:	Rectum Abdomen						
Circle severity:	No pain 0 1 2	3 4 5 6 7 8	9 10 Worst pai	n			

Pelvic Pain Symptoms: None

□ My pelvic symptoms <u>affect my ability</u> to, OR <u>I feel worse when I try</u> to: Check all that apply.

□ Sleep	Bend forward		Climb stairs	Derform work duties		
Bathe	□Squat down		□ Sitting tolerance	□Recreational activities		
Get dressed	Lift items		□ Standing tolerance	□Social events		
Wear tight clothing	Reach overhead		Walking distance	□Travel		
Cook meals	Get out of bed		Drive a car	Exercise for health		
Do housework	□Stand up from a c	hair	□ Run errands/shop	Do Kegel exercises		
Do yard work	Get out of a car					
PenisLow back	 Scrotum Buttocks Side/waist Urethra 	□ Groin □ Bladder □ Tailbone 4 5 6 7	 Over tight surgical sca Sides of hips Back of hips 9 10 Worst Pain 	ars		
□ I limit the <i>frequency</i> of sexual activity because of my pain.			 I have to <i>interrupt</i> intercourse due to pain. I <i>avoid it</i> altogether due to my pain. Last intercourse attempt: 			
The pain occurs during: During erection During ejaculation						
For how long? 🛛 Only during sexual activity 📮 For a few hours afterwards 📮 For a day or more afterwards						

Location: \Box Penis/scrotum \Box Deep in my pelvis \Box In my back

Circle pain severity: No pain $0_1_2_3_4_5_6_7_9_10$ Worst Pain



PELVIC FLOOR INFORMED CONSENT FOR EVALUATION AND TREATMENT OF PELVIC FLOOR DYSFUNCTIONS

The term, "informed consent," means that the potential risks, benefits and alternative of therapy evaluation and treatment have been explained to me. The therapist provides a wide range of services, and I understand that I will receive information at the initial visit concerning the evaluation, treatment and options available for my condition.

I understand that if I am referred to physical therapy for pelvic floor dysfunction, it may be beneficial for my therapist to perform a muscle assessment of the pelvic floor, initially and periodically to assess muscle strength, length and range of motion and scar mobility. Palpation of these muscles is most direct and accessible if done via the vagina and/or rectum. Pelvic floor dysfunctions include pelvic pain syndromes, urinary incontinence, fecal incontinence, dyspareunia or pain with intercourse, pain from episiotomy or scarring, vulvodynia, vestibulitis, persistent sacroiliac joint dysfunction/ low back pain, or other similar complications. Evaluation of my condition may include observation, soft tissue mobilization, use of vaginal cones, and vaginal or rectal sensors for muscle education and/or electrical stimulation.

I understand that the benefits of the vaginal/rectal assessment have been explained to me. I understand that if I am uncomfortable with the assessment or treatment procedures AT ANY TIME, I will inform my therapist, and the procedure will be discontinued and an alternative discussed with me.

Treatment procedures for pelvic floor dysfunctions may include, without limitation, education, exercise, stimulation, ultrasound, and several manual therapy techniques including massage, joint and soft tissue mobilization. The therapist will explain all of the treatment procedures to me, and I may choose to not participate with all or part of the treatment plan. I understand that no guarantees have been or can be provided to me regarding the success of therapy.

I have received instructions on a three-dimensional, anatomical model to better understand muscle and organ location, hand placement, and palpation techniques that are common in the treatment of pelvic floor muscle conditions or disorders.

Potential Risks: I may experience an increase in my current level of pain or discomfort, or an aggravation of my existing injury. This discomfort is usually temporary. If pain or discomfort does not subside in 1 to 3 days, I agree to contact my therapist.

I have read, or had read to me, the foregoing, and any questions which may have occurred to me have been answered to my satisfaction. I understand the risks, benefits and alternatives of the treatment.

Based on the information I have received from the therapist, I voluntarily agree to standard assessment and muscular treatment techniques of the perineal area.

____ I am comfortable with only the therapist performing the evaluation in the room.

____ I would prefer to have a chaperone in the room while the therapist performs the evaluation.



*** If you are pregnant, have an infection of any kind, have vaginal dryness, are less than 6 weeks post partum or post surgery, have severe pelvic pain, sensitivity to creams, please inform the therapist prior to pelvic floor assessment.

Evaluation:

Patient Signature	Date	Therapist Signature	Date
Treatments:			
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date
Patient Signature	Date	Therapist Signature	Date

Taken from APTA Sect on Women's Health, Sher Pelvic Health and Healing and 3rd Source not named from the internet.

https://goconfluent-my.sharepoint.com/personal/mkaye_golcr_com/Documents/MaryKaye/FORMS/PATIENT/Originals #2 2020/LCR PF/PF informed consent LCR 1-20.docx